



**You have downloaded a document from
RE-BUS
repository of the University of Silesia in Katowice**

Title: Two new genera and species of the Gigantometopini (Hemiptera, Heteroptera, Miridae, Isometopinae) from Borneo with remarks on the distribution of the tribe

Author: Artur Taszakowski, Junggon Kim, Claas Damken, Rodzay A. Wahab, Aleksander Herczek, Sunghoon Jung

Citation style: Taszakowski Artur, Kim Junggon, Damken Claas, Wahab Rodzay A., Herczek Aleksander, Jung Sunghoon. (2020). Two new genera and species of the Gigantometopini (Hemiptera, Heteroptera, Miridae, Isometopinae) from Borneo with remarks on the distribution of the tribe. "ZooKeys" (Vol. 941 (2020), s. 71-89), doi 10.3897/zookeys.941.47432



Uznanie autorstwa - Licencja ta pozwala na kopiowanie, zmienianie, rozprowadzanie, przedstawianie i wykonywanie utworu jedynie pod warunkiem oznaczenia autorstwa.



UNIwersYTET ŚLĄSKI
W KATOWICACH



Biblioteka
Uniwersytetu Śląskiego



Ministerstwo Nauki
i Szkolnictwa Wyższego

Two new genera and species of the Gigantometopini (Hemiptera, Heteroptera, Miridae, Isometopinae) from Borneo with remarks on the distribution of the tribe

Artur Tazakowski^{1*}, Junggon Kim^{2*}, Claas Damken³, Rodzay A. Wahab³,
Aleksander Herczek¹, Sunghoon Jung^{2,4}

1 Institute of Biology, Biotechnology and Environmental Protection, Faculty of Natural Sciences, University of Silesia in Katowice, Bankowa 9, 40-007 Katowice, Poland **2** Laboratory of Systematic Entomology, Department of Applied Biology, College of Agriculture and Life Sciences, Chungnam National University, Daejeon, South Korea **3** Institute for Biodiversity and Environmental Research, Universiti Brunei Darussalam, Jalan Universiti, BE1410, Darussalam, Brunei **4** Department of Smart Agriculture Systems, College of Agriculture and Life Sciences, Chungnam National University, Daejeon, South Korea

Corresponding author: Artur Tazakowski (artur.tazakowski@us.edu.pl); Sunghoon Jung (jung@cnu.ac.kr)

Academic editor: F. Konstantinov | Received 21 October 2019 | Accepted 2 May 2020 | Published 16 June 2020

<http://zoobank.org/B3C9A4BA-B098-4D73-A60C-240051C72124>

Citation: Tazakowski A, Kim J, Damken C, Wahab RA, Herczek A, Jung S (2020) Two new genera and species of the Gigantometopini (Hemiptera, Heteroptera, Miridae, Isometopinae) from Borneo with remarks on the distribution of the tribe. ZooKeys 941: 71–89. <https://doi.org/10.3897/zookeys.941.47432>

Abstract

Two new genera, each represented by a single new species, *Planicapitus luteus* Tazakowski, Kim & Herczek, **gen. et sp. nov.** and *Bruneimetopus simulans* Tazakowski, Kim & Herczek, **gen. et sp. nov.**, are described from Borneo. Detailed photographs of male habitus and genital structures are presented. The checklist with distributional records for all known taxa of Gigantometopini is also provided.

Keywords

Asia, biodiversity, distribution, jumping tree bugs, plant bugs, true bugs

* Contributed equally as the first authors.

Introduction

The Isometopinae are a highly autapomorphic group possessing paired ocelli which are absent in all other members of the plant bug family Miridae (Herczek 1993, Namyatova and Cassis 2016, Yasunaga et al. 2017). This subfamily was considered as the sister group to all other subfamilies based on morphology (Schuh 1974, 1976), but recent works using molecular data do not support this hypothesis (Schuh et al. 2009, Jung and Lee 2012). Therefore, additional works are needed to understand phylogenetic position of this subfamily.

The group has a worldwide distribution, but the majority of known taxa are thermophilic, and occur in the tropics, subtropics, and warm temperate climate zones (Eyles 1971, Schuh 2002–2013, Cassis and Schuh 2012, Namyatova and Cassis 2016, Yasunaga et al. 2017). Due to scarce information on habits, biology, and food preference, the representatives are relatively rare in collections with many species known from singletons or only a handful of specimens (Eyles 1971, Namyatova and Cassis 2016, Herczek et al. 2018). Currently, six tribes, 45 genera and more than 250 species of Isometopinae are known (Namyatova and Cassis 2016, Yasunaga et al. 2016, 2017, Krüger 2018, Herczek et al. 2018) of which 19 species are fossil taxa (Herczek 1993, Schuh 2002–2013, Herczek and Popov 2012, 2014). The Isometopini and Myiommini are the most species-rich isometopine tribes known worldwide (Namyatova and Cassis 2016). The Electromyiommini is an extinct tribe and contains four genera and 14 species; all were described from Baltic amber (Herczek 1993). The Diphlebini includes only a single genus, *Diphleps* Bergroth, 1924 (Schuh 2002–2013). Yasunaga et al. (2017) created the new tribe Sophianini comprising two genera previously classified within Myiommini: *Alcecoris* McAtee & Malloch, 1924 and *Sophianus* Distant, 1904. Sophianini includes ten species (Yasunaga et al. 2017).

The last tribe is Gigantometopini created by Herczek (1993) to accommodate a single species *Gigantometopus rossi* Schwartz & Schuh, 1990. In 2002 *Isometopidea gryllocephala* Miyamoto, Yasunaga & Hayashi, 1996 was transferred to a newly created genus *Astroscometopus* Yasunaga & Hayashi and its inclusion to the tribe Gigantometopini was suggested (Yasunaga and Hayashi 2002). In 2004 another species of *Isometopidea* was described, *Isometopidea formosana* Lin, and the next year it was transferred to the genus *Astroscometopus* (Lin 2005). The same year Yasunaga (2005) described a new species, representing a new genus, *Kohnometopus fraxini* Yasunaga, 2005 in the tribe Myiommini Bergroth, 1924. Next, Akingbohunge (2012) described the second representative of the genus *Gigantometopus*, *G. schuhi* Akingbohunge, 2012. Yasunaga et al. (2017) transferred *Isometopidea yangi* (Lin 2005) to the genus *Kohnometopus*, suggested that this genus seemed better placed in Gigantometopini rather than in Myiommini, and also proposed to place the genus *Isometopidea* Poppius, 1913 (with the single species *Isometopidea lieweni* Poppius, 1913) in the tribe Gigantometopini. Moreover, it was found that the identity of the specimen of *I. lieweni* from Taiwan (Lin and Yang 2004) was based on a misidentification and it is a representative of an undescribed species. Subsequently Herczek et al. (2018) described one more genus and species within Gigantometopini, *Sulawesimetopus henryi* Herczek, Gorczyca & Taszakowski.

The most characteristic feature of Gigantometopini distinguishing it from other tribes is the large numbers of trichobothria (five or six on both mesofemur and metafemur) (Yasunaga et al. 2017).

In this paper, two new genera and species *Planicapitus luteus* gen. et sp. nov. and *Bruneimetopus simulans* gen. et sp. nov. are diagnosed and described; photographic images of habitus and genital structures, as well as scanning electron micrographs of the selected structures of both species are provided.

Materials and methods

The specimens were imaged by the following equipment: Leica M205C stereo microscope with high diffuse dome illumination Leica LED5000 HDI, Leica DFC495 digital camera and Leica application suite 4.9.0 software; Leica DM 3000 upright light microscope with Leica MC 190 HD digital camera and Leica Application Suite 4.12.0 software. SEM photographs were obtained using Phenom XL field emission scanning electron microscope at 5 and 10 kV accelerating voltage with a BackScatter Detector (BSD). Graphic editor Adobe Photoshop CS6 was used to prepare the figures. In case of legs, the preparations for SEM were made with methods traditionally used in morphological studies (e.g. Kanturski et al. 2015, Herczek et al. 2018). In contrast, during preparation of other photographs, steps that can damage the specimen e.g., washing, dehydration and sputter-coating with a film of electrically conducting material, have not been applied. Specimens on original glue boards were only cleaned with a brush and mounted on aluminium stubs with double-sided adhesive carbon tape. Next, the specimens were covered with anti-static spray.

Map was prepared in SAGA GIS 7.1.1 (<http://www.saga-gis.org>) using WGS84 datum and EPSG: 3395 (World Mercator cylindrical projection).

Measurements were made with Leica application suite 4.9.0 software and are presented in millimetres (mm). Terminology of morphological structures mainly follows Herczek et al. (2018) and Kim and Jung (2019). Dissections of male genitalia were performed using Kerzhner and Konstantinov's (1999) technique. The terminology for genital structures follows Konstantinov (2003). The study was based on material deposited in the collection of the Royal Belgian Institute of Natural Sciences (RBINS) and material recently collected by Claas Damken during an extensive survey of the Heteroptera fauna of Brunei Darussalam, deposited at Universiti Brunei Darussalam Museum, Brunei Darussalam (UBDM). From 2013 to 2015, sampling took place at different locations and forest types across the Bornean Sultanate using a range of methods (e.g., generator-powered light traps, sweep netting, collecting by hand, litter sifting, pitfall traps, Malaise traps, examination of bycatch from other studies). During this field survey, more than 400 species of Heteroptera were collected, including several species of Isometopinae (<https://tinyurl.com/Brunei-Isometopinae>).

Taxonomy

Planicapitus Tazsakowski, Kim & Herczek, gen. nov.

<http://zoobank.org/E38884A2-CDD1-4C4F-95A5-E8AA1F979AA5>

Type species. *Planicapitus luteus* Tazsakowski, Kim & Herczek, sp. nov.

Diagnosis. Distinguished by vertical, flattened head, not punctured but wrinkled and distinctly higher than wide, dorsally extending to level of highest point of pronotum; vertex convex, protruding above eye level; width of vertex slightly larger than eye width; dorsum and pleurites of thorax with deep and dense punctures; calli slightly marked, tarsi two segmented, claw without subapical tooth; labium reaching third abdominal segment; right paramere very small, short, dagger-shaped; left paramere ca. 2.5 times as long as right one.

Description. Male. Body oval, slightly elongated (Fig. 1A). Head clearly higher than wide, dorsally extending to highest point of pronotum, flattened, impunctate but wrinkled; Antenna thin (particularly segments III and IV). Labium reaching third abdominal segment (Fig. 1B). Pronotal collar with row of punctures. Pronotum distinctly punctate, distinctly carinate at sides, with slightly upturned lateral margins; calli slightly marked, separated by shallow fossa. Scutellum convex, wider than long, basomedially clearly depressed. Thoracic pleura distinctly punctate (Fig. 1C). Ostiolar peritreme small, strongly convex and covered with fine spines (Fig. 4A–C). Mesofemora with five trichobothria (Fig. 3C, D). Tarsi two segmented, claws without apical tooth (Fig. 3 E–G). Genitalia: genital capsule trapeziform, with two longitudinal sutures at sides (Fig. 5A, B); aedeagus delicate, membranous, with weakly sclerotized dorsal wall of phallosome, endosoma sacciform and membranous, weakly sclerotized (Fig. 5A, B, E). Left paramere scythe-shaped, sensory lobe with several long setae, apical process elongate (Fig. 5A–C); right paramere very small, short, dagger-shaped (Fig. 5A, B, D).

Remarks. Affiliation of *Planicapitus luteus* to the Gigantometopini is clearly confirmed by the following features: compound eyes relatively small, significantly separated from each other, pronotum deeply punctate and elongate, calli separated by shallow fossa, pronotal collar demarcated by row of punctures, inflated scutellum, and five mesofemoral trichobothria (Herczek 1993, Yasunaga et al. 2017).

Set of features mentioned in the diagnosis clearly differ the new genus from other genera belonging to Gigantometopini. *Planicapitus luteus* belongs to the smallest representatives of tribe. The new genus is similar in size to *Isometopidea lieweni* which body length of the only known specimen equals to 3.0 mm. It is a female, so probably (like other representatives of tribe) males reach a smaller body size (Poppius 1913, Yasunaga 2005, Herczek et al. 2018). *Isometopidea* further differs from newly described genus by the structure of the head, which is not higher than wide, somewhat rounded and not strongly flattened in front. *Sulawesimetopus*, the second comparatively small-sized genus of the Gigantometopini, is slightly larger and can be distinguished from the new genus by the three segmented tarsi and punctured head. Other representatives of Gigantometopini are a way larger than the new genus in body size.




Figure 1. *Planicapitus luteus*: dorsal (A), ventral (B) and lateral (C) views.

Etymology. Combined from Latin adjective: *planus*, flat and noun: *caput*, *capitis*, head; gender masculine.

***Planicapitus luteus* Tazakowski, Kim & Herczek, sp. nov.**

<http://zoobank.org/F66DC352-A83C-4661-897A-D101551CDC2C>

Figures 1–5

Diagnosis. See generic diagnosis.

Description. Male. Body shiny, yellow-brownish, covered by semi-erect pale-brown seta (Fig. 1A–C). **Head:** yellow-whitish, 1.5 times as high as wide, compound eyes reddish yellow, vertex orange, convex, 1.3 times as wide as eye width in dorsal view. Frons whitish, with two small dark brown spots ventrally extending into large Y-shaped brown macula; gena whitish yellow (Figs 1C, 2A, B). Antenna yellowish. Labium shiny, yellowish, segment IV with brown apex (Fig. 1B). **Thorax:** pronotum yellow, semi-transparent laterally; exposed part of mesoscutum yellow, scutellum yellowish brown, with apex white and lateral angles narrowly whitish, 0.7 as long as wide. Pleura yellowish brown, with red stripe from propleuron to episternum (Fig. 1B, C). Ostiolar peritreme ivory, evaporative area yellow (Fig. 1B, C). Claval commissure 0.6 times as long as scutellum. **Hemelytron:** in various shades of yellow, median part with




Figure 2. *Planicapitus luteus*: head in frontal view (A), lateral view of head (B), left antenna (C).




Figure 3. *Planicapitus luteus*: head in dorsal view (**A**, **B**), femur of middle leg in ventrolateral view, showing trichobothrial pattern (**C**, **D**), pretarsus of foreleg, lateral view (**E**), tarsus of middle leg, lateral view (**F**), pretarsus of middle leg, ventral view (**G**).

two whitish spots. Cuneus 0.9 times as long as wide, yellowish, with white spot in basal inner corner. Membrane pale grey, semi-transparent, with two cells. **Legs:** coxae pale, almost white, femora yellow-white (Fig. 1B, C), with brown spots, tibiae yellow with




Figure 4. Scent gland, evaporatory area (**A, B**) and peritreme (**C**) of *Planicapitus luteus*.




Figure 5. Male genitalia of *Planicapitus luteus*: genital capsule in dorsal (**A**) and caudal (**B**) views, left paramere (**C**), right paramere (**D**), aedeagus in dorsal view (**E**).

dark brown spots, tarsi yellow (Fig. 1B, C). **Abdomen:** bicoloured, dark brown, except for pale yellow genital segment (Fig. 1B, C). **Genitalia:** as described above. **Measurements:** given in the Table 1.

Etymology. From Latin adjective *luteus*, yellow.

Biology. Unknown.

Material examined. *Holotype* (♂): 'BORNEO, Sabah / DANUM VALLEY / 70km W Lahad Datu / M.J. & J.P. Duffels // East Ridge Trail / 150m / 2.XII.1989 // sample Sab. 53 / understorey rainforest, / at light'. The holotype is deposited in RBINS.

***Bruneimetopus* Tazsakowski, Kim & Herczek, gen. nov.**

<http://zoobank.org/63C3A9E7-12A1-4CB7-AB92-6F702663401D>

Type species. *Bruneimetopus simulans* Tazsakowski, Kim & Herczek, sp. nov.

Diagnosis. Distinguished by vertical, slightly flattened head, not punctured but wrinkled and higher than wide, dorsally not extending to level of highest point of pronotum; vertex slightly convex, protruding above eye level, width of vertex equal to eye width; dorsum and pleurites of thorax with deep and dense punctures; calli slightly marked, tarsi two segmented, claw with very small, barely noticeable apical tooth; labium reaching third abdominal segment, right paramere well developed, with knee-shaped sensory lobe; left paramere ca. 1.5 times as long as right one.

Description. Male. Body oval, slightly elongate (Fig. 6A). Head higher than wide, dorsally not extending to highest point of pronotum, slightly flattened, impunctate but wrinkled. Antenna thin (particularly segments III and IV). Labium reaching third abdominal segment. Pronotal collar with row of punctures posteriorly. Pronotum distinctly punctate, calli slightly marked, separated by shallow fossa. Scutellum slightly convex, baso-medially clearly depressed. Thoracic pleura distinctly punctate. Ostiolar peritreme small, moderately convex and covered with very fine spines (Fig. 8). Mesofemora with five, metafemora with six trichobothria (Fig. 7F, G). Tarsi two segmented, claws with very small, barely noticeable apical tooth (Fig. 7D, E). Genitalia: genital capsule trapeziform (Fig. 9A), aedeagus delicate, endosoma sacciform and membranous, very weakly sclerotized inside, outer subapical and apical part more sclerotic, clothed with dense spinules (Fig. 9A, D). Left paramere scythe-shaped, sensory lobe with several long setae, apical process elongated (Fig. 9A, B); right paramere left paramere ca. 1.5 times as long as right one, with knee-shaped sensory lobe (Fig. 9C).

Remarks. Affiliation of *Bruneimetopus* to the Gigantometopini is clearly confirmed by the same set of features as for *Planicapitus* (see above). It is also indicated by presence of six metafemoral trichobothria (the specimen of *Planicapitus luteus* is devoid of hindlegs).

As in the case of *Planicapitus*, set of features mentioned in the diagnosis clearly differ the new genus from other genera belonging to Gigantometopini. The newly described genera are very similar morphologically to each other. However, in addition to small differences in the proportions of body parts and coloration, they can easily be distinguished by the completely different shape and size of the right paramere. This was a premise to describe them in separate genera.

Etymology. Name combines Brunei (the type locality) with part of the generic name *Isometopus*, the type genus of the subfamily.

***Bruneimetopus simulans* Tazsakowski, Kim & Herczek, sp. nov.**

<http://zoobank.org/40B3C1AB-7913-4AA4-9AE1-82B098AE5B23>

Figures 6–9

Diagnosis. See generic diagnosis.

Description. Male. Body shiny, in various shades of yellow and brown, covered by semi-erect pale brown and brown setae (Fig. 6A–C). **Head:** brownish yellow, 1.4




Figure 6. *Bruneimetopus simulans*: holotype: dorsal (A), ventral (B) and lateral (C) views.

times as high as wide, compound eyes reddish, vertex white, slightly convex, as wide as eye in dorsal view. Frons dark brown between eyes, yellowish below inferior margin of eyes; clypeus brown; gena yellow (Fig. 7A–C). Antenna yellowish, segments III and IV




Figure 7. *Bruneimetopus similans*: holotype: head in frontal view (A), lateral view of head (B), head in dorsal view (C), tarsus of middle leg, lateral view (D), pretarsus of middle leg, lateral view (E), paratype: femur of middle leg in ventral view, showing trichobothrial pattern (F), femur of hind leg in ventral view, showing trichobothrial pattern (G).

darker. Labium shiny, yellowish, segment IV brown (Fig. 6B). **Thorax:** pronotum dark yellow, lateral margins semi-transparent and slightly raised, slightly wider at front; posterior margin whitish. Exposed part of mesoscutum brown with yellow tinge. Scutel-




Figure 8. *Bruneimetopus simulans* (holotype) scent gland, evaporatory area (A, B) and peritreme (C).




Figure 9. Male genitalia of *Bruneimetopus simulans*: holotype: genital capsule in dorsal view (A), left paramere (B), right paramere (C), aedeagus (D).

lum dark brown, with white apical part and black extreme apex, 0.6 times as long as wide, covered by semi-erect setae. Propleuron dark yellow, meso- and metapleurons dark brown with dark yellow tinge. Ostiolar peritreme ivory, evaporative area yellow-brown (Fig. 4D–F). Claval commissure comparatively long, 0.5 times as long as scutellum. **Hemelytron:** in various yellow and brown shades: median, posterior part and cuneus in 2/3 of their length semi-transparent, whitish yellow, base of hemelytra and clavus yellow-brown, part neighbouring with cuneal fracture and 1/3 length of cuneus dark brown. Cuneus 0.9 as long as wide. Membrane pale grey, semi-transparent, with two cells. **Legs:** coxae yellowish pale, femora yellow-white, with brown spots apically, tibiae yellow with four or five dark brown, irregular rings, tarsi yellow (Fig. 6A, B). **Abdomen:** bicolored: first two segments yellowish to brown, others dark brown (Fig. 6B, C). **Genitalia:** as described above. **Measurements:** given in the Table 1.

Table 1. Comparison of metric features of *Planicapitus luteus* and *Bruneimetopus simulans*.

	<i>P. luteus</i>	<i>B. simulans</i> holotype	<i>B. simulans</i> paratype	<i>B. simulans</i> average
Body length	2.61	2.52	2.47	2.50
Body width	1.24	1.13	–	1.13
Head length	0.22	0.19	0.19	0.19
Head width	0.58	0.51	0.52	0.52
Head height	0.86	0.71	0.68	0.71
Dorsal width of eye	0.20	0.17	0.19	0.18
Vertex width	0.26	0.17	0.19	0.18
Antennal segments I:II:III:IV	0.10:0.62:0.67:0.21	0.10:0.59:0.78:0.20	0.09:0.60:0.82:0.20	0.10:0.60:0.80:0.20
Labium length	1.30	1.26	–	1.26
Labial segments I:II:III:IV	0.35:0.26:0.30:0.37	0.34:0.36:0.23:0.39	–	0.34:0.36:0.23:0.39
Pronotum length	0.52	0.48	0.43	0.46
Anterior width of pronotum	0.51	0.48	0.44	0.46
Posterior width of pronotum	1.19	1.07	1.07	1.07
Mesoscutum length	0.10	0.12	0.10	0.11
Scutellum length	0.42	0.42	0.49	0.46
Scutellum width	0.61	0.68	0.59	0.64
Claval commissure	0.27	0.23	–	0.23
1 st femur length	0.73	0.64	0.68	0.66
1 st tibia length	0.76	0.73	0.76	0.75
1 st length of tarsus	0.26	0.20		
1 st length of tarsus I: II	0.09:0.22	0.08:0.17	0.08:0.17	0.08:0.17
2 nd femur length	0.86	0.73	0.73	0.73
2 nd tibia length	0.96	0.84	0.88	0.86
2 nd length of tarsus	0.21	0.19	0.19	0.19
2 nd length of tarsus I: II	0.08:0.18	0.08:0.16	0.07:0.15	0.08:0.16
3 rd femur length	–	–	0.92	0.92
3 rd femur width	–	–	0.23	0.23
3 rd tibia length	–	–	1.28	1.28
3 rd tarsus length	–	–	0.22	0.22
3 rd length of tarsus I: II	–	–	0.09:0.15	0.09:0.15
Heme length	1.99	1.84	–	1.84
Corium length	1.55	1.25	–	1.25
Cuneus length	0.29	0.23	–	0.23
Cuneus width	0.33	0.27	–	0.27

Etymology. The species name *simulans* (resembling) is the present participle of the Latin verb *simulo* (to make like or to assume the appearance of anything), in allusion to the resemblance of this species to *Planicapitus luteus*.

Biology. Unknown. Two specimens were collected in a mangrove forest (Fig. 10) by a Malaise trap, together with several other specimens of Isometopinae.

Material examined. *Holotype* (♂): 'Borneo, BRUNEI, Tutong // Tutong area, mangroves forest / small stream near water edge, Malaise / trap 1; 16.viii.2014, leg: C. Dam-



Figure 10. Malaise trap at the collecting site of holotype of *Bruneimetopus simulans*.

ken / 4°46'9.54"N, 114°36'20.64"E // ID code: tutong.mangroves.01780'; **Paratype** (♂): 'Borneo, BRUNEI, Labu FR. / mangrove forest, Malaise trap ID4 / 06.viii.2018; leg: C. Damken / 4°50'53.11"N, 117°7'45.65"E// ID code: labu.mangroves.01777'. The holotype and paratype are deposited in the UBDM.

Distributional remarks

In total, only 49 imagines of Gigantometopini representing eleven species were ever recorded. Four species are known only from the holotype: *Gigantometopus rossi*, *Gigantometopus schuhi*, *Isometopidea lieweni*, and *Planicapitus luteus*. Below we present the complete checklist of Gigantometopini with distributional records (Fig. 11) and biological information (following Poppius 1913, Schwartz and Schuh 1990, Miyamoto et al. 1996, Yasunaga and Hayashi 1996, Lin and Yang 2004, Yasunaga 2005, Akingbohunge 2012, Yasunaga et al. 2017, Herczek et al. 2018):

Gigantometopini Herczek, 1993

Astroscometopus Yasunaga & Hayashi, 2002

***Astroscometopus formosanus* (Lin, 2004)**

Isometopidea formosana Lin, 2004




Figure 11. Distribution of Gigantometopini: **A** *I. lieweni* **B** *G. rossi* **C** *G. schuhi* and *B. simulans* **D** *P. luteus* **E** *S. henryi* **F** *A. formosanus* **G** *K. yangi* **H** *A. formosanus* and *I. lieweni* nec **I** *A. gryllocephalus* **J** *A. gryllocephalus* and *K. fraxini*

1♂, Taiwan, Nantou, Chunyang (Fig. 11H), 11 Jun–9 Jul 2002, malaise trap

1♂, Taiwan, Pingtung, Hengchun, Kenting National Park (Fig. 11F), 10 Mar–14 Apr 2005, malaise trap

***Astroscometopus gryllocephalus* (Miyamoto, Yasunaga, & Hayashi, 1996)**

Isometopidea gryllocephala Miyamoto, Yasunaga, & Hayashi, 1996

1♀, Japan, Ryukyu Arc., Ishigaki Is., Shiramizu (Fig. 11J), 19 Mar 1993, sweeping, grasses growing on the subtropical jungle floor near a montane stream.

1♀, Japan, Ryukyu Arc., Ishigaki Is., Mt. Yarabudake (Fig. 11J), 10 Mar 1999, the bark of the subtropical ash, *Fraxinus griffithii*

1♂, 8 final instar nymphs, Japan, Ryukyu Arc., Ishigaki Is., Mt. Fukami-Omoto (Fig. 11J), 18 Mar 2000, the bark of the subtropical ash, *Fraxinus griffithii*

1♀, Japan, Ryukyu Arc., Iriomote Is. (Fig. 11I), 2 Mar 2002, root of an unidentified broadleaved tree

Gigantometopus Schwartz & Schuh, 1990

***Gigantometopus rossi* Schwartz & Schuh, 1990**

1♀, Indonesia, Sumatra, Sumatera Barat, Mangani, mine near Kota Tinggi, 700 m a.s.l. (Fig. 11B), 20 Jul 1983

***Gigantometopus* cf. *rossi* Schwartz & Schuh, 1990**

1♀, South India

***Gigantometopus schuhi* Akingbohungbe, 2012**

1♂, Brunei, Borneo, Bukit Sulang near Lamunin (Fig. 11C), 20 Aug–10 Sep 1982, insecticide fogging on *Shorea macrocarpa*

Isometopidea Poppius, 1913

***Isometopidea lieweni* Poppius, 1913**

1♀, Sri Lanka, Anuradhapura (Fig. 11A), 21 Dec

***Isometopidea lieweni* nec Poppius, 1913**

1♀, Taiwan, Nantou, Lienhachi (Fig. 11H), Nov 1984, malaise trap

Kohnometopus Yasunaga, 2005

***Kohnometopus fraxini* Yasunaga, 2005**

1♂, Japan, Ryukyu Arc., Ishigaki Is., Mt. Fukami-Omoto (Fig. 11J), 28 Sep 2002; 6♀♀, 22 May 2002

1♀, Japan, Ryukyu Arc., Ishigaki Is., Mt. Yarabudake (Fig. 11J), 1 Jun 2002; 2♂♂, 6♀♀, 28 Nov 2002; 1♀, 2 Oct 2002; all specimens of this species were collected on two trees of *Fraxinus griffithii*

***Kohnometopus yangi* (Lin, 2005)**

Isometopidea yangi Lin 2005

1♂, Taiwan, Taitung, Peinan Panchiu Station (Fig. 11G), 19 Nov–16 Dec 2004, malaise trap; 2♂♂, 2♀♀, 19 Nov–16 Dec 2004, malaise trap; 1♀, 7 Oct–19 Nov 2004, malaise trap; 3♂♂, 1♀, 16 Dec 2004–17 Feb 2005, malaise trap

Sulawesimetopus Herczek, Gorczyca & Tazsakowski, 2018

***Sulawesimetopus henryi* Herczek, Gorczyca & Tazsakowski, 2018**

3♂♂, Indonesia, Sulawesi Utara (Fig. 11E), 8–18 Nov 1985

5♂♂, 1♀, Indonesia, Sulawesi Utara, Dumonga-Bone National Park, Hogg's Back Subcamp, 660 m. a.s.l. (Fig. 11E), 15 Nov 1985

Planicapitus Tazsakowski, Kim & Herczek, gen. nov.

***Planicapitus luteus* Tazsakowski, Kim & Herczek, sp. nov.**

1♂, Malaysia, Borneo, Sabah Danum Valley, East Ridge Trail, 150 m. a.s.l., (Fig. 11D), 2 Dec 1989, understorey rainforest, at light

Bruneimetopus Tazsakowski, Kim & Herczek, gen. nov.

***Bruneimetopus simulans* Tazsakowski, Kim & Herczek, sp. nov.**

1♂, Brunei, Borneo, Tutong area (Fig. 11C), 16 Aug 2014, mangroves forest (Fig. 10), malaise trap

1♂, Brunei, Borneo, Labu FR. (Fig. 11C), 6 Aug 2018, mangroves forest, malaise trap

Acknowledgements

We are greatly indebted to Dr Jerome Constant from the Royal Belgian Institute of Natural Sciences, Prof Dominik Chłond (University of Silesia in Katowice) for a loan and the assistance with the loan material and Dr Agnieszka Bugaj-Nawrocka (University of Silesia in Katowice) for help in making the map. Claas Damken gratefully acknowledges the support by Ulmar Grafe (Universiti Brunei Darussalam) during the

fieldwork and research on the Heteroptera of Brunei Darussalam. The authors also thank Brunei Forestry Department and Ministry of Primary Resources and Tourism for permission to conduct field work in Brunei Darussalam. The Biodiversity Research and Innovation Centre provided export permits. We also want to thank the reviewers as well as the editors for all the comments that improved this manuscript.

We wanted to apologize to Dr Anna Namyatova for misquoting the work she co-authored in our latest publication. Correct title: Namyatova AA, Cassis G (2016) Review of the seven new species of Isometopinae (Heteroptera: Miridae) in Australia and discussion of distribution and host plant associations of the subfamily on the worldwide basis. *Austral Entomology* 55: 392–422. <https://doi.org/10.1111/aen.12202>

This work was supported by Korea Environment Industry & Technology Institute (KEITI) through Exotic Invasive Species Management Program, funded by Korea Ministry of Environment (MOE)(2018002270005).

The research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Education (2018R1A6A3A01013374), and was funded through the UBD postdoctoral fellowship (Research and export permit file no: UBD/CAN–387(b)(SAA); UBD/ADM/R3(z)Pt.; UBD/PNC2/2/RG/1(293)).

References

- Akingbohunge AE (2012) A note on *Gigantometopus* Schwartz and Schuh (Heteroptera: Miridae: Isometopinae) with the description of a new species from Borneo. *Entomologica Americana* 118(1/4): 130–132. <https://doi.org/10.1664/12-RA-015.1>
- Cassis G, Schuh RT (2012) Systematics, biodiversity, biogeography, and host associations of the Miridae (Insecta: Hemiptera: Heteroptera: Cimicomorpha). *Annual Review of Entomology* 57: 377–404. <https://doi.org/10.1146/annurev-ento-121510-133533>
- Eyles AC (1971) List of Isometopidae (Heteroptera: Cimicoidea). *New Zealand Journal of Science* 14, 940–944.
- Herczek A, Popov YA (2012) A new peculiar isometopine genus (Hemiptera: Heteroptera: Miridae) from the Eocene Baltic amber. *Zootaxa* 3196: 64–68. <https://doi.org/10.11646/zootaxa.3196.1.4>
- Herczek A (1993) Systematic position of Isometopinae Fieb. (Miridae, Heteroptera) and their intrarelationships. *Prace Naukowe Uniwersytetu Śląskiego, Katowice*, 1357: 1–86.
- Herczek A, Gorczyca J, Taszakowski A (2018) *Sulawesimetopus henryi*, a new genus and species of Isometopinae (Hemiptera, Heteroptera, Miridae) from Sulawesi. In: Wheeler Jr AG (Ed.) *A Festschrift Recognizing Thomas J. Henry for a Lifetime of Contributions to Heteropteran Systematics*. *ZooKeys* 796: 147–161. <https://doi.org/10.3897/zookeys.796.21273>
- Jung S, Lee S (2012) Molecular phylogeny of the plant bugs (Heteroptera: Miridae) and the evolution of feeding habits. *Cladistics* 28: 50–79. <https://doi.org/10.1111/j.1096-0031.2011.00365.x>

- Kanturski M, Karcz J, Wiczorek K (2015) Morphology of the European species of the aphid genus *Eulachnus* (Hemiptera: Aphididae: Lachninae) – a SEM comparative and integrative study. *Micron* 76: 23–36. <https://doi.org/10.1016/j.micron.2015.05.004>
- Kerzhner IM, Konstantinov FV (1999) Structure of the aedeagus in Miridae (Heteroptera) and its bearing to suprageneric classification. *Acta Societatis Zoologicae Bohemicae* 63: 117–137.
- Konstantinov FV (2003) Male genitalia in Miridae (Heteroptera) and their significance for suprageneric classification of the family. Part I: general review, Isometopinae and Psallopinae. *Belgian Journal of Entomology* 5: 3–36.
- Kim J, Jung S (2019) Phylogeny of the plant bug subfamily Mirinae (Hemiptera: Heteroptera: Cimicomorpha: Miridae) based on total evidence analysis. *Systematic Entomology* 44, 686–698. <https://doi.org/10.1111/syen.12348>
- Lin CS (2004) Seven new species of Isometopinae (Hemiptera: Miridae) from Taiwan. *Formosan Entomologist* 24: 317–326.
- Lin CS (2005) New or little-known Isometopinae from Taiwan (Hemiptera: Miridae). *Formosan Entomologist* 25: 195–201.
- Lin CS, Yang CT (2004) Isometopinae (Hemiptera: Miridae) from Taiwan. *Formosan Entomologist* 24: 27–42.
- Miyamoto S, Yasunaga T, Hayashi M (1996) Description of a new isometopine plant bug, *Isometopidea gryllocephala*, found on Ishigaki Island, Japan (Insecta, Heteroptera, Miridae). *Species Diversity* 1: 107–110. <https://doi.org/10.12782/specdiv.1.107>
- Namyatova AA, Cassis G (2016) Review of the seven new species of Isometopinae (Heteroptera: Miridae) in Australia and discussion of distribution and host plant associations of the subfamily on a worldwide basis. *Austral Entomology* 55: 392–422. <https://doi.org/10.1111/aen.12202>
- Poppius B (1913) Zur Kenntniss der Miriden, Isometopiden, Anthocoriden, Nabiden und Schizopteriden Ceylon's. *Entomologisk Tidskrift* 34: 239–260. <https://doi.org/10.5962/bhl.part.1634>
- SAGA Development Team (2019) System for Automated Geoscientific Analyses (SAGA) (Version 7.1.1) Institute of Geography at the University of Hamburg, Germany. <http://www.saga-gis.org> [Accessed: 6 February 2019]
- Schuh RT (1974) The Orthotylinae and Phylinae (Hemiptera: Miridae) of South Africa with a phylogenetic analysis of the antmimetic tribes of the two subfamilies for the world. *Entomologica Americana* 47: 1–332.
- Schuh RT (1976) Pretarsal structure in the Miridae (Hemiptera) with a cladistic analysis of relationships within the family. *American Museum Novitates* 2601: 1–39.
- Schuh RT (2002–2013) On-line systematic catalog of plant bugs (Insecta: Heteroptera: Miridae). <http://research.amnh.org/pbi/catalog> [Accessed 19 February 2019]
- Schuh RT, Weirauch C, Wheeler WC (2009) Phylogenetic relationships within the Cimicomorpha (Hemiptera: Heteroptera): a total-evidence analysis. *Systematic Entomology* 34: 15–48. <https://doi.org/10.1111/j.1365-3113.2008.00436.x>
- Schwartz MD, Schuh RT (1990) The world's largest Isometopine, *Gigantometopus rossi*, new genus and new species (Heteroptera: Miridae). *Journal New York Entomological Society* 98(1): 9–13.

- Yasunaga T (2005) Isometopinae plant bugs (Heteroptera: Miridae) preferably inhabiting *Fraxinus griffithii* on Ishigaki Island of the Ryukyus, Japan. Tijdschrift voor Entomologie 148: 341–349. <https://doi.org/10.1163/22119434-900000179>
- Yasunaga T, Yamada K, Tsai JF (2017) Taxonomic review of the plant bug subfamily Isometopinae for Taiwan and Japanese Southwest Islands, with descriptions of new taxa (Hemiptera: Heteroptera: Miridae: Isometopinae). Zootaxa 4365(4): 421–439
- Yasunaga T, Duangthisan J, Yamada K, Artchawakom T (2016) Further records of the plant bug subfamily Isometopinae from Thailand (Heteroptera: Miridae) with description of three new species. Tijdschrift voor Entomologie 159: 89–96. <https://doi.org/10.1163/22119434-15902003>
- Yasunaga T, Hayashi M (2002) New or little known isometopine plant bugs from Japan (Heteroptera: Miridae). Tijdschrift voor Entomologie 145: 95–101. <https://doi.org/10.1163/22119434-900000103>